

Wie Mythen der traumafokussierten Psychotherapie eine adäquate Versorgung erschweren

Ein Plädoyer zur Implementierung evidenzbasierter Verfahren in Deutschland

Philipp Herzog, Tim Kaiser & Ad de Jongh

Literatur

- Bachmann, C. J., Czwikla, J., Jacobs, H., Fegert, J. M., & Hoffmann, F. (2021). Prävalenz und Versorgung der Posttraumatischen Belastungsstörung in Deutschland: Eine bundesweite Auswertung von Krankenkassendaten aus den Jahren 2008 und 2017. *Psychiatrische Praxis*, 48 (6), 316–323. <https://doi.org/10.1055/a-1347-5410>
- Bækkelund, H., Karlsrud, I., Hoffart, A., & Arnevik, E. A. (2021). Stabilizing group treatment for childhood-abuse related PTSD: A randomized controlled trial. *European Journal of Psychotraumatology*, 12 (1), 1859079. <https://doi.org/10.1080/20008198.2020.1859079>
- Bisson, J. I., Roberts, N. P., Andrew, M., Cooper, R., & Lewis, C. (2013). Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *Cochrane Database of Systematic Reviews*, 12. <https://doi.org/10.1002/14651858.CD003388.pub4>
- Bohus, M., Kleindienst, N., Hahn, C., Müller-Engelmann, M., Ludäscher, P., Steil, R., Fydrich, T., Kuehner, C., Resick, P. A., Stiglmayr, C., Schmahl, C., & Priebe, K. (2020). Dialectical Behavior Therapy for Posttraumatic Stress Disorder (DBT-PTSD) Compared with Cognitive Processing Therapy (CPT) in Complex Presentations of PTSD in Women Survivors of Childhood Abuse: A Randomized Clinical Trial. *JAMA Psychiatry*, 77 (12), 1235–1245. <https://doi.org/10.1001/jamapsychiatry.2020.2148>
- Bongaerts, H., Voorendonk, E. M., van Minnen, A., & de Jongh, A. (2021). Safety and effectiveness of intensive treatment for complex PTSD delivered via home-based telehealth. *European Journal of Psychotraumatology*, 12 (1), 1860346. <https://doi.org/10.1080/20008198.2020.1860346>
- Bongaerts, H., Voorendonk, E. M., Van Minnen, A., Rozendaal, L., Telkamp, B. S. D., & de Jongh, A. (2022). Fully remote intensive trauma-focused treatment for PTSD and Complex PTSD. *European Journal of Psychotraumatology*, 13 (2), 2103287. <https://doi.org/10.1080/20008066.2022.2103287>
- Boterhoven De Haan, K. L., Lee, C. W., Fassbinder, E., Van Es, S. M., Menninga, S., Meewisse, M. L. et al. (2020). Imagery rescripting and eye movement desensitisation and reprocessing as treatment for adults with post-traumatic stress disorder from childhood trauma: Randomised clinical trial. *British Journal of Psychiatry*, 217 (5), 609–615. <https://doi.org/10.1192/bjp.2020.158>
- Bredemeier, K., Larsen, S., Shivakumar, G., Grubbs, K., McLean, C., Tress, C. et al. (2022). A comparison of prolonged exposure therapy, pharmacotherapy, and their combination for PTSD: What works best and for whom; study protocol for a randomized trial. *Contemporary Clinical Trials*, 119, 106850. <https://doi.org/10.1016/j.cct.2022.106850>
- Cloitre, M. (2016). Commentary on De Jongh Et Al. (2016) Critique of Istss Complex Ptsd Guidelines: Finding the Way Forward. *Depression and Anxiety*, 33 (5), 355–356. <https://doi.org/10.1002/da.22493>
- Cloitre, M., Koenen, K. C., Cohen, L. R., & Han, H. (2002). Skills training in affective and interpersonal regulation followed by exposure: A phase-based treatment for PTSD related to childhood abuse. *Journal of Consulting and Clinical Psychology*, 70 (5), 1067–1074. <https://doi.org/10.1037//0022-006x.70.5.1067>

- Cusack, K., Jonas, D. E., Forneris, C. A., Wines, C., Sonis, J., Middleton, J. C. et al. (2016). Psychological treatments for adults with posttraumatic stress disorder: A systematic review and meta-analysis. *Clinical Psychology Review*, 43 (290), 128–141. <https://doi.org/10.1016/j.cpr.2015.10.003>
- De Jongh, A., Resick, P. A., Zoellner, L. A., van Minnen, A., Lee, C. W., Monson, C. M. et al. (2016). Critical analysis of the current treatment guidelines for complex PTSD in adults. *Depression and Anxiety*, 33 (5), 359–369. <https://doi.org/10.1002/da.22469>
- Deisenhofer, A.-K., Delgadillo, J., Rubel, J. A., Böhnke, J. R., Zimmermann, D., Schwartz, B., & Lutz, W. (2018). Individual treatment selection for patients with posttraumatic stress disorder. *Depression and Anxiety*, 35 (6), 541–550. <https://doi.org/10.1002/da.22755>
- Ehlers, A. (1999). Posttraumatische Belastungsstörung. Göttingen: Hogrefe.
- Ehlers, A., Grey, N., Wild, J., Stott, R., Liness, S., Deale, A. et al. (2013). Implementation of Cognitive Therapy for PTSD in routine clinical care: Effectiveness and moderators of outcome in a consecutive sample. *Behaviour Research and Therapy*, 51 (11), 742–752. <https://doi.org/10.1016/j.brat.2013.08.006>
- Ehlers, A., Hackmann, A., Grey, N., Wild, J., Liness, S., Albert, I. et al. (2014). A randomized controlled trial of 7-day intensive and standard weekly cognitive therapy for PTSD and emotion-focused supportive therapy. *American Journal of Psychiatry*, 171 (3), 294–304. <https://doi.org/10.1176/appi.ajp.2013.13040552>
- Ehring, T., Welboren, R., Morina, N., Wicherts, J. M., Freitag, J., & Emmelkamp, P. M. G. (2014). Meta-analysis of psychological treatments for posttraumatic stress disorder in adult survivors of childhood abuse. *Clinical Psychology Review*, 34 (8), 645–657. <https://doi.org/10.1016/j.cpr.2014.10.004>
- Epping, J., Zwaan, M. de, & Geyer, S. (2018). Gesünder nach der Psychotherapie? Sekundärdatenanalyse der Arbeitsunfähigkeitszeiten vor und nach ambulanter Verhaltenstherapie, tiefenpsychologisch fundierter und analytischer Psychotherapie. *PPmP - Psychotherapie · Psychosomatik · Medizinische Psychologie*, 68 (8), 337–345. <https://doi.org/10.1055/s-0043-120346>
- Fischer, G., Reddemann, L., Barwinski-Fäh, R., & Bering, R. (2003). Traumaadaptierte tiefenpsychologisch fundierte und analytische Psychotherapie. *Psychotherapeut*, 48 (3), 199–209. <https://doi.org/10.1007/s00278-003-0302-9>
- Foa, E. B., Bredemeier, K., Acieno, R., Rosenfield, D., Muzzy, W., Tuerk, P. W., Zandberg, L. J. et al. (2022). The efficacy of 90-min versus 60-min sessions of prolonged exposure for PTSD: A randomized controlled trial in active-duty military personnel. *Journal of Consulting and Clinical Psychology*, 90 (6), 503–512. <https://doi.org/10.1037/ccp0000739>
- Foa, E. B., Hembree, E. A., Rothbaum, B. O., Kierdorf, T., & Höhr, H. (2014). Handbuch der Prolongierten Exposition: Basiskonzepte und Anwendung - eine Anleitung für Therapeuten. Lichtenau: G.P. Probst Verlag.
- Foa, E. B., McLean, C. P., Zang, Y., Rosenfield, D., Yadin, E., Yarvis, J. S. et al. (2018). Effect of Prolonged Exposure Therapy Delivered Over 2 Weeks vs 8 Weeks vs Present-Centered Therapy on PTSD Symptom Severity in Military Personnel A Randomized Clinical Trial. *JAMA - Journal of the American Medical Association*, 319 (4), 354–364. <https://doi.org/10.1001/jama.2017.21242>
- Gahleitner, S. B. (2020). »Ich gehe heute so weit, zu meinen, dass wir unsere Worte auf die Goldwaage des Mitgefühls legen sollten«: Ein Interview mit Luise Reddemann, geführt von Silke Birgitta Gahleitner. *Trauma & Gewalt*, 14 (3), 244–247. <https://doi.org/10.21706/tg-14-3-244>

- Gillespie, K., Duffy, M., Hackmann, A., & Clark, D. M. (2002). Community based cognitive therapy in the treatment of posttraumatic stress disorder following the Omagh bomb. *Behaviour Research and Therapy*, 40 (4), 345–357. [https://doi.org/10.1016/s0005-7967\(02\)00004-9](https://doi.org/10.1016/s0005-7967(02)00004-9)
- Halvorsen, J. Ø., Stenmark, H., Neuner, F., & Nordahl, H. M. (2014). Does dissociation moderate treatment outcomes of narrative exposure therapy for PTSD? A secondary analysis from a randomized controlled clinical trial. *Behaviour Research and Therapy*, 57 (1), 21–28. <https://doi.org/10.1016/j.brat.2014.03.010>
- Herzog, P., & Kaiser, T. (2022). Is it worth it to personalize the treatment of PTSD? – A variance-ratio meta-analysis and estimation of treatment effect heterogeneity in RCTs of PTSD. *Journal of Anxiety Disorders*, 102611. <https://doi.org/10.1016/j.janxdis.2022.102611>
- Herzog, P., Voderholzer, U., Gärtner, T., Osen, B., Svitak, M., Doerr, R. et al. (2020). Predictors of outcome during inpatient psychotherapy for posttraumatic stress disorder: A single-treatment, multi-site, practice-based study. *Psychotherapy Research*. <https://doi.org/10.1080/10503307.2020.1802081>
- Hoeboer, C. M., De Kleine, R. A., Molendijk, M. L., Schoorl, M., Oprel, D. a. C., Mouthaan, J. et al. (2020). Impact of dissociation on the effectiveness of psychotherapy for post-traumatic stress disorder: Meta-analysis. *BJPsych Open*, 6 (3), e53. <https://doi.org/10.1192/bjo.2020.30>
- Jayawickreme, N., Cahill, S. P., Riggs, D. S., Rauch, S. A. M., Resick, P. A., Rothbaum, B. O., & Foa, E. B. (2014). Primum non nocere (first do no harm): Symptom worsening and improvement in female assault victims after prolonged exposure for PTSD. *Depression and Anxiety*, 31 (5), 412–419. <https://doi.org/10.1002/da.22225>
- Kahn, G. (2008). Das Innere-Kinder-Retten – eine Methode der imaginativen Traumatherapie. *Trauma & Gewalt*, 2 (1), 48–52.
- Kolthof, K. A., Voorendonk, E. M., Van Minnen, A., & De Jongh, A. (2022). Effects of intensive trauma-focused treatment of individuals with both post-traumatic stress disorder and borderline personality disorder. *European Journal of Psychotraumatology*, 13 (2), 2143076. <https://doi.org/10.1080/20008066.2022.2143076>
- König, J., Resick, P. A., Karl, R., & Rosner, R. (2012). Posttraumatische Belastungsstörung: Ein Manual zur Cognitive Processing Therapy (12012th ed.). Göttingen: Hogrefe.
- Kuester, A., Niemeyer, H., & Knaevelsrud, C. (2016). Internet-based interventions for posttraumatic stress: A meta-analysis of randomized controlled trials. *Clinical Psychology Review*, 43, 1–16. <https://doi.org/10.1016/j.cpr.2015.11.004>
- Lampe, A., Mitmansgruber, H., Gast, U., Schüßler, G., & Reddemann, L. (2008). Therapieevaluation der Psychodynamisch Imaginativen Traumatherapie (PITT) im stationären Setting. *Neuropsychiatrie*. <https://doi.org/10.5414/NEPBand22189>
- Levitt, J. T., & Cloitre, M. (2005). A Clinician's Guide to STAIR/MPE: Treatment for PTSD Related to Childhood Abuse. *Cognitive and Behavioral Practice*, 12, 40–52. [https://doi.org/10.1016/S1077-7229\(05\)80038-0](https://doi.org/10.1016/S1077-7229(05)80038-0)
- Lewis, C., Roberts, N. P., Andrew, M., Starling, E., & Bisson, J. I. (2020). Psychological therapies for post-traumatic stress disorder in adults: Systematic review and meta-analysis. *European Journal of Psychotraumatology*, 11 (1). <https://doi.org/10.1080/20008198.2020.1729633>
- Lindauer, M., Linden, M., & Muschalla, B. (2022). Verständnis ambulanter Verhaltenstherapeuten von Belastungsreaktionen und PTBS. *Verhaltenstherapie*, 32 (4), 186–195. <https://doi.org/10.1159/000526482>

- McLean, C. P., Levy, H. C., Miller, M. L., & Tolin, D. F. (2022). Exposure therapy for PTSD: A meta-analysis. *Clinical Psychology Review*, 91, 102115. <https://doi.org/10.1016/j.cpr.2021.102115>
- Meehl, P. E. (1997). Credentialled persons, credentialled knowledge. *Clinical Psychology: Science and Practice*, 4(2), 91–98. <https://doi.org/10.1111/j.1468-2850.1997.tb00103.x>
- Murray, H., Grey, N., Warnock-Parkes, E., Kerr, A., Wild, J., Clark, D. M., & Ehlers, A. (2022). Ten misconceptions about trauma-focused CBT for PTSD. *The Cognitive Behaviour Therapist*, 15, e33. <https://doi.org/10.1017/S1754470X22000307>
- National Institute for Clinical Excellence (NICE). (2018). Post-Traumatic Stress Disorder (National Institute for Clinical Excellence (UK) & National Collaborating Centre for Mental Health (UK), Ed.; Issue 26). Published by Gaskell and the British Psychological Society. <https://doi.org/10.1192/pb.30.9.357-a>
- Orel, D. A. C., Hoeboer, C. M., Schoorl, M., de Kleine, R. A., Cloitre, M., Wigard, I. G. et al. (2021). Effect of Prolonged Exposure, intensified Prolonged Exposure and STAIR+Prolonged Exposure in patients with PTSD related to childhood abuse: A randomized controlled trial. *European Journal of Psychotraumatology*, 12 (1), 1851511. <https://doi.org/10.1080/20008198.2020.1851511>
- Pawils, S., Nick, S., Metzner, F., Lotzin, A., & Schäfer, I. (2017). Versorgungssituation von Kindern, Jugendlichen und Erwachsenen mit sexuellen Gewalterfahrungen in Deutschland: Ein kritischer Überblick. *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz*, 60 (9), 1046–1054. <https://doi.org/10.1007/s00103-017-2601-2>
- Peterson, A. L., Blount, T. H., Foa, E. B., Brown, L. A., McLean, C. P., Mintz, J. et al. (Consortium to Alleviate PTSD). (2023). Massed vs Intensive Outpatient Prolonged Exposure for Combat-Related Posttraumatic Stress Disorder: A Randomized Clinical Trial. *JAMA Network Open*, 6 (1), e2249422. <https://doi.org/10.1001/jamanetworkopen.2022.49422>
- Pittig, A., Kotter, R., & Hoyer, J. (2019). The Struggle of Behavioral Therapists With Exposure: Self-Reported Practicability, Negative Beliefs, and Therapist Distress About Exposure-Based Interventions. *Behavior Therapy*, 50 (2), 353–366. <https://doi.org/10.1016/j.beth.2018.07.003>
- Raabe, S., Ehring, T., Marquenie, L., Arntz, A., & Kindt, M. (2022). Imagery Rescripting as a stand-alone treatment for posttraumatic stress disorder related to childhood abuse: A randomized controlled trial. *Journal of Behavior Therapy and Experimental Psychiatry*, 77, 101769. <https://doi.org/10.1016/j.jbtep.2022.101769>
- Reddemann, L. (2003). Psychotherapie von Suchterkrankungen als Traumafolgestörungen. *PiD - Psychotherapie im Dialog*, 4 (2), 136–139. <https://doi.org/10.1055/s-2003-39521>
- Reddemann, L., & Piedfort-Marin, O. (2017). Stabilization in the treatment of complex post-traumatic stress disorders: Concepts and principles. *European Journal of Trauma & Dissociation*, 1 (1), 11–17. <https://doi.org/10.1016/j.ejtd.2017.01.009>
- Rosner, R., Henkel, C., Ginkel, K., & Mestel, R. (2010). Was passiert nach der stationären Stabilisierung mit komplex traumatisierten PTB-Patientinnen? *Zeitschrift Für Psychiatrie, Psychologie und Psychotherapie*, 58 (2), 127–135. <https://doi.org/10.1024/1661-4747/a000017>
- Sack, M. (2020). Schonende Traumatherapie: Ressourcenorientierte Behandlung von Traumafolgestörungen (2. Aufl.). Stuttgart: Schattauer.
- Schäfer, I., Gast, U., Hofmann, A., Knaevelsrud, C., Lampe, A., Liebermann, P. et al. (2019). S3-Leitlinie Posttraumatische Belastungsstörung. Berlin/Heidelberg: Springer.
- Schauer, M., Elbert, T., & Neuner, F. (2011). Narrative exposure therapy: A short-term treatment for traumatic stress disorders. Göttingen: Hogrefe.

- Schnurr, P. P., Chard, K. M., Ruzek, J. I., Chow, B. K., Resick, P. A., Foa, E. B. et al. (2022). Comparison of Prolonged Exposure vs Cognitive Processing Therapy for Treatment of Posttraumatic Stress Disorder Among US Veterans. *JAMA Network Open*, 5 (1), e2136921. <https://doi.org/10.1001/jamanetworkopen.2021.36921>
- Schock, K., Rosner, R., Wenk-Ansohn, M., & Knaevelsrud, C. (2010). Retraumatisierung – Annäherung an eine Begriffsbestimmung. *PPmP - Psychotherapie · Psychosomatik · Medizinische Psychologie*, 60 (07), 243–249. <https://doi.org/10.1055/s-0030-1248268>
- Schwartz, B., Cohen, Z. D., Rubel, J. A., Zimmermann, D., Wittmann, W. W., & Lutz, W. (2020). Personalized treatment selection in routine care: Integrating machine learning and statistical algorithms to recommend cognitive behavioral or psychodynamic therapy. *Psychotherapy Research*, 31 (1), 33–51. <https://doi.org/10.1080/10503307.2020.1769219>
- Shapiro, F., & Maxfield, L. (2002). Eye movement desensitization and reprocessing (EMDR): Information processing in the treatment of trauma. *Journal of Clinical Psychology*, 58 (8), 933–946. <https://doi.org/10.1002/jclp.10068>
- Shearing, V., Lee, D., & Clohessy, S. (2011). How do clients experience reliving as part of trauma-focused cognitive behavioural therapy for posttraumatic stress disorder? *Psychology and Psychotherapy*, 84 (4), 458–475. <https://doi.org/10.1111/j.2044-8341.2010.02012.x>
- Sloan, D. M., Marx, B. P., Lee, D. J., & Resick, P. A. (2018). A brief exposure-based treatment vs cognitive processing therapy for posttraumatic stress disorder: A randomized noninferiority clinical trial. *JAMA Psychiatry*, 75 (3), 233–239. <https://doi.org/10.1001/jamapsychiatry.2017.4249>
- Sloan, D. M., Marx, B. P., Resick, P. A., Young-McCaughan, S., Dondanville, K. A., Straud, C. L. et al. (2022). Effect of Written Exposure Therapy vs Cognitive Processing Therapy on Increasing Treatment Efficiency Among Military Service Members With Posttraumatic Stress Disorder. *JAMA Network Open*, 5 (1), e2140911. <https://doi.org/10.1001/jamanetworkopen.2021.40911>
- Smucker, Mervin., Reschke, Konrad., & Kögel, Betty. (2008). Imagery rescripting & reprocessing therapy: Behandlungsmanual für Typ I Trauma. Düren: Shaker.
- Sommer, J. (2016). Die psychotherapeutische Versorgungsrealität komplex traumatisierter Menschen in Deutschland. *Trauma & Gewalt*, 10 (4), 308–319. <https://doi.org/10.21706/tg-10-4-308>
- Stirman, S. W., Cohen, Z. D., Lunney, C. A., DeRubeis, R. J., Wiley, J. F., & Schnurr, P. P. (2021). A personalized index to inform selection of a trauma-focused or non-trauma-focused treatment for PTSD. *Behaviour Research and Therapy*, 142, 103872. <https://doi.org/10.1016/j.brat.2021.103872>
- van Minnen, A., Hendriks, L., & Olff, M. (2010). When do trauma experts choose exposure therapy for PTSD patients? A controlled study of therapist and patient factors. *Behaviour Research and Therapy*, 48 (4), 312–320. <https://doi.org/10.1016/j.brat.2009.12.003>
- van Pelt, Y., Fokkema, P., de Roos, C., & de Jongh, A. (2021). Effectiveness of an intensive treatment programme combining prolonged exposure and EMDR therapy for adolescents suffering from severe post-traumatic stress disorder. *European Journal of Psychotraumatology*, 12 (1), 1917876. <https://doi.org/10.1080/20008198.2021.1917876>
- van Vliet, N. I., Huntjens, R. J. C., van Dijk, M. K., Bachrach, N., Meewisse, M.-L., & de Jongh, A. (2021a). Phase-based treatment versus immediate trauma-focused treatment for post-traumatic stress disorder due to childhood abuse: Randomised clinical trial. *BJPsych Open*, 7 (6), e211. <https://doi.org/10.1192/bjo.2021.1057>
- van Vliet, N. I., Huntjens, R. J. C., van Dijk, M. K., Bachrach, N., Meewisse, M.-L., & de Jongh, A. (2021b). Phase-based treatment versus immediate trauma-focused treatment for post-traumatic

stress disorder due to childhood abuse: Randomised clinical trial. *BJPsych Open*, 7 (6), e211. <https://doi.org/10.1192/bjo.2021.1057>

van Woudenberg, C., Voorendonk, E. M., Bongaerts, H., Zoet, H. A., Verhagen, M., Lee, C. W. et al. (2018). Effectiveness of an intensive treatment programme combining prolonged exposure and eye movement desensitization and reprocessing for severe post-traumatic stress disorder. *European Journal of Psychotraumatology*, 9 (1), 1487225. <https://doi.org/10.1080/20008198.2018.1487225>

Voorendonk, E.M., De Jongh, A., Roozendaal, L. &, Van Minnen, A. (2020). Trauma-focused treatment outcome for Complex PTSD patients: Results of an intensive treatment programme. *European Journal of Psychotraumatology*, 11 (1), 1783955. <https://doi.org/10.1080/20008198.2020.1783955>

Wagenmans, A., Van Minnen, A., Slijpen, M. &, De Jongh, A. (2018). The Impact of childhood sexual abuse on the outcome of intensive trauma-focused Treatment for PTSD. *European Journal of Psychotraumatology*, 9 (1), 1430962. <https://doi.org/10.1080/20008198.2018.1430962>

Zoet, H. A., De Jongh, A. & van Minnen, A. (2021). Somatoform dissociative symptoms have no impact on the outcome of trauma-focused treatment for severe PTSD. *Journal of Clinical Medicine*, 10, 1553. <https://doi.org/10.3390/jcm10081553>