

## Körperliches Training: eine additive Behandlungsmethode?

Viola Oertel, Miriam Bieber, Daniela Schmidt, Esra Görgülü & Kirsten Zabel

Ainsworth, B. E., Haskell, W. L., Leon, A. S., Jacobs, J. D., Montoye, H. J., Sallis, J. F. et al. (1993). Compendium of physical activities: classification of energy costs of human physical activities. *Medicine and science in sports and exercise*, 25(1), 71–80.

Banzer, W. (Hrsg.). (2017). *Körperliche Aktivität und Gesundheit*. Berlin: Springer. <https://doi.org/10.1007/978-3-662-50335-5>

Bressington, D., Badnapurkar, A., Inoue, S., Yeung Ma, H., Chien, W.T., Nelson, D. & Gray, R. (2018). Physical Health Care for People with Severe Mental Illness: the Attitudes, Practices, and Training Needs of Nurses in Three Asian Countries. *Int J Environ Res Public Health*. 15(2): 343.

Broocks, A., Bandelow, B., Pekrun, G., George, A., Meyer, T., Bartmann, U. et al. (1998). Comparison of aerobic exercise, clomipramine, and placebo in the treatment of panic disorder. *The American Journal of Psychiatry*, 155(5), 603–609. <https://doi.org/10.1176/ajp.155.5.603>

Broocks, A. & Sommer, M. (2005). Sport und Psyche. *Standards der Sportmedizin, Deutsche Zeitschrift für Sportmedizin*, Jahrgang 56, Nr. 11.

Cless, K. & Matura, S. (2017). Psychische Gesundheit. In V. Oertel & S. Matura (Hrsg.), *Bewegung und Sport gegen Burnout, Depressionen und Ängste* (S. 13–17). Berlin: Springer. [https://doi.org/10.1007/978-3-662-53938-5\\_2](https://doi.org/10.1007/978-3-662-53938-5_2)

Conn, V. S. (2010). Depressive symptom outcomes of physical activity interventions: meta-analysis findings. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine*, 39(2), 128–138. <https://doi.org/10.1007/s12160-010-9172-x>

Convertino, V. A. (2007). Blood volume response to physical activity and inactivity. *The American Journal of the Medical Sciences*, 334(1), 72–79. <https://doi.org/10.1097/MAJ.0b013e318063c6e4>

Cotman, C. W., Berchtold, N. C. & Christie, L.-A. (2007). Exercise builds brain health: key roles of growth factor cascades and inflammation. *Trends in Neurosciences*, 30(9), 464–472. <https://doi.org/10.1016/j.tins.2007.06.011>

DGPPN & KBV (Hrsg.) (2015). *S3-Leitlinie/Nationale VersorgungsLeitlinie Unipolare Depression – Langfassung*, 2. Auflage. Version 5, 238.

Dunn, A. L., Trivedi, M. H. & O’Neal, H. A. (2001). Physical activity dose-response effects on outcomes of depression and anxiety. *Medicine and Science in Sports and Exercise*, 33(6 Suppl), S587-597; discussion 609-610.

Eggermont, L., Swaab, D., Luiten, P. & Scherder, E. (2006). Exercise, cognition and Alzheimer's disease: more is not necessarily better. *Neuroscience and Biobehavioral Reviews*, 30(4), 562–575. <https://doi.org/10.1016/j.neubiorev.2005.10.004>

Firth, J., Carney, R., Elliott, R., French, P., Parker, S., McIntyre, R. et al. (2018). Exercise as an intervention for first-episode psychosis: a feasibility study: Exercise intervention for first-episode psychosis. *Early Intervention in Psychiatry*, 12(3), 307–315. <https://doi.org/10.1111/eip.12329>

Firth, J., Cotter, J., Carney, R. & Yung, A. R. (2017). The pro-cognitive mechanisms of physical exercise in people with schizophrenia: Pro-cognitive mechanisms of exercise in schizophrenia. *British Journal of Pharmacology*, 174(19), 3161–3172. <https://doi.org/10.1111/bph.13772>

Firth, J., Rosenbaum, S., Stubbs, B., Gorczynski, P., Yung, A. R. & Vancampfort, D. (2016). Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. *Psychological Medicine*, 46(14), 2869–2881. <https://doi.org/10.1017/S0033291716001732>

Fogarty, M. & Happell, B. (2005). Exploring the benefits of an exercise program for people with schizophrenia: a qualitative study. *Issues in Mental Health Nursing*, 26(3), 341–351. <https://doi.org/10.1080/01612840590915711>

Fuchs, D. R. & Göhner, D. W. (o. J.). Hoch motiviert! Für mehr Sport und Bewegung – das Trainingsbuch zur Broschüre, 65.

Göhner, W. & Fuchs, R. (2006). Änderung des Gesundheitsverhaltens: MoVo-Gruppenprogramme für körperliche Aktivität und gesunde Ernährung. Göttingen: Hogrefe.

Gorczynski, P. & Faulkner, G. (2010). Exercise therapy for schizophrenia. (Cochrane Schizophrenia Group, Hrsg.), *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD004412.pub2>

Gordon, E. M., Laumann, T. O., Gilmore, A. W., Newbold, D. J., Greene, D. J., Berg, J. J. et al. (2017). Precision Functional Mapping of Individual Human Brains. *Neuron*, 95(4), 791-807.e7. <https://doi.org/10.1016/j.neuron.2017.07.011>

Görgülü, E., Bieber, M., Engeroff, T., Schmidt, D., Zabel, K., Oertel, V., Reif, A. (in Vorbereitung). Physical activity, physical self-perception and depression symptoms in patients with major depressive disorder: A mediation analysis.

Heggelund, J., Nilsberg, G. E., Hoff, J., Morken, G. & Helgerud, J. (2011). Effects of high aerobic intensity training in patients with schizophrenia: a controlled trial. *Nordic Journal of Psychiatry*, 65(4), 269–275. <https://doi.org/10.3109/08039488.2011.560278>

Hoffman, B. M., Babyak, M. A., Craighead, W. E., Sherwood, A., Doraiswamy, P. M., Coons, M. J. et al. (2011). Exercise and pharmacotherapy in patients with major depression: one-year follow-up of the SMILE study. *Psychosomatic Medicine*, 73(2), 127–133. <https://doi.org/10.1097/PSY.0b013e31820433a5>

Kaluza, G. (2011). *Stressbewältigung*. Berlin: Springer. <https://doi.org/10.1007/978-3-642-13720-4>

Keller-Varady, K., Falkai, P. & Malchow, B. (2017). Sporttherapie bei Schizophrenie. *PSYCH up2date*, 11(01), 51–65. <https://doi.org/10.1055/s-0042-118429>

Knuttgen, H. (2008). Basic Exercise Physiology (S. 3–16). <https://doi.org/10.1002/9780470693766.ch1>

Krogh, J., Nordentoft, M., Sterne, J. A. C. & Lawlor, D. A. (2011). The effect of exercise in clinically depressed adults: systematic review and meta-analysis of randomized controlled trials. *The Journal of Clinical Psychiatry*, 72(4), 529–538. <https://doi.org/10.4088/JCP.08r04913blu>

Laursen, T. M. (2011). Life expectancy among persons with schizophrenia or bipolar affective disorder. *Schizophrenia Research*, 131(1–3), 101–104. <https://doi.org/10.1016/j.schres.2011.06.008>

Loprinzi, P. D. & Wade, B. (2019). Exercise and cardiorespiratory fitness on subjective memory complaints. *Psychology, Health & Medicine*, 24(6), 749–756. <https://doi.org/10.1080/13548506.2018.1557713>

Loprinzi, P.D., Loenneke, J.P., Ahmed, H.M., Blaha, M.J., (2016). Joint effects of objectively-measured sedentary time and physical activity on all-cause mortality. *Preventive Medicine*. 90: 47.

Mehnert, A., Veers, S., Howaldt, D., Braumann, K.M., Koch, U., Schulz, K.H., (2011). Effects of a physical exercise rehabilitation group program on anxiety, depression, body image, and health-related quality of life among breast cancer patients. *Onkologie*;34(5):248-53.

Melo, M. C. A., Daher, E. D. F., Albuquerque, S. G. C. & de Bruin, V. M. S. (2016). Exercise in bipolar patients: A systematic review. *Journal of Affective Disorders*, 198, 32–38. <https://doi.org/10.1016/j.jad.2016.03.004>

Mittal, V. A., Vargas, T., Osborne, K. J., Dean, D., Gupta, T., Ristanovic, I. et al. (2017). Exercise Treatments for Psychosis: A Review. *Current Treatment Options in Psychiatry*, 4(2), 152–166. <https://doi.org/10.1007/s40501-017-0112-2>

Montgomery, S. A. & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *The British Journal of Psychiatry: The Journal of Mental Science*, 134, 382–389.

National Institute for Health and Care Excellence (2013): Physical activity: brief advice for adults in primary care. In: NICE public health guidance 44. Manchester.

Oeland, A.M., Laessoe, U., Olesen, A.V., Munk-Jørgensen, P., (2010). Impact of exercise on patients with depression and anxiety. *Nord J Psychiatry*. May 4;64(3):210-7

Oertel, V. & Matura, S. (Hrsg.). (2017). *Bewegung und Sport gegen Burnout, Depressionen und Ängste*. Berlin: Springer. <https://doi.org/10.1007/978-3-662-53938-5>

Oertel-Knöchel, V. & Hänsel, F. (2015). *Aktiv für die Psyche: Sport und Bewegungsinterventionen bei psychisch kranken Menschen*. Heidelberg: Springer-Verlag.

Oertel-Knöchel, V., Mehler, P., Thiel, C., Steinbrecher, K., Malchow, B., Tesky, V. et al. (2014). Effects of aerobic exercise on cognitive performance and individual psychopathology in depressive and schizophrenia patients. *European Archives of Psychiatry and Clinical Neuroscience*, 264(7), 589–604. <https://doi.org/10.1007/s00406-014-0485-9>

Osborn, D. P. J., Nazareth, I. & King, M. B. (2007). Physical activity, dietary habits and Coronary Heart Disease risk factor knowledge amongst people with severe mental illness: a cross sectional comparative study in primary care. *Social Psychiatry and Psychiatric Epidemiology*, 42(10), 787–793. <https://doi.org/10.1007/s00127-007-0247-3>

Pajonk, F.-G., Wobrock, T., Gruber, O., Scherk, H., Berner, D., Kaizl, I. et al. (2010). Hippocampal plasticity in response to exercise in schizophrenia. *Archives of General Psychiatry*, 67(2), 133–143. <https://doi.org/10.1001/archgenpsychiatry.2009.193>

Paton, C., Esop, R., Young, C. & Taylor, D. (2004). Obesity, dyslipidaemias and smoking in an inpatient population treated with antipsychotic drugs. *Acta Psychiatrica Scandinavica*, 110(4), 299–305. <https://doi.org/10.1111/j.1600-0447.2004.00372.x>

Petruzzello, S. J., Landers, D. M., Hatfield, B. D., Kubitz, K. A. & Salazar, W. (1991). A meta-analysis on the anxiety-reducing effects of acute and chronic exercise. Outcomes and mechanisms. *Sports Medicine (Auckland, N.Z.)*, 11(3), 143–182. <https://doi.org/10.2165/00007256-199111030-00002>

Poulin, M.-J., Chaput, J.-P., Simard, V., Vincent, P., Bernier, J., Gauthier, Y. et al. (2007). Management of antipsychotic-induced weight gain: prospective naturalistic study of the effectiveness of a supervised exercise programme. *The Australian and New Zealand Journal of Psychiatry*, 41(12), 980–989. <https://doi.org/10.1080/00048670701689428>

van Praag, H., Shubert, T., Zhao, C. & Gage, F. H. (2005). Exercise enhances learning and hippocampal neurogenesis in aged mice. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 25(38), 8680–8685. <https://doi.org/10.1523/JNEUROSCI.1731-05.2005>

Rethorst, C. D., Wipfli, B. M. & Landers, D. M. (2009). The antidepressive effects of exercise: a meta-analysis of randomized trials. *Sports Medicine (Auckland, N.Z.)*, 39(6), 491–511. <https://doi.org/10.2165/00007256-200939060-00004>

Rethorst, C.D., Trivedi, M. (2013). Evidence-Based Recommendations for the Prescription of Exercise for Major Depressive Disorder; *Journal of Psychiatric Practice*. 19(3):204–212, MAY 2013. <https://doi.org/10.1097/01.pra.0000430504.16952.3e>

Roick, C., Fritz-Wieacker, A., Matschinger, H., Heider, D., Schindler, J., Riedel-Heller, S. et al. (2007). Health habits of patients with schizophrenia. *Social Psychiatry and Psychiatric Epidemiology*, 42(4), 268–276. <https://doi.org/10.1007/s00127-007-0164-5>

Rovio, S., Kåreholt, I., Helkala, E.-L., Viitanen, M., Winblad, B., Tuomilehto, J. et al. (2005). Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *The Lancet. Neurology*, 4(11), 705–711. [https://doi.org/10.1016/S1474-4422\(05\)70198-8](https://doi.org/10.1016/S1474-4422(05)70198-8)

Schuch, F., Vancampfort, D., Firth, J., Rosenbaum, S., Ward, P., Reichert, T. et al. (2017). Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. *Journal of Affective Disorders*, 210, 139–150. <https://doi.org/10.1016/j.jad.2016.10.050>

Schulz, K.-H., Meyer, A. & Langguth, N. (2012). Körperliche Aktivität und psychische Gesundheit. *Bundesgesbl*; 55: 55-65

Siems, M. (2018). Effekte von positivem Feedback auf die Zielerreichung bei einer bewegungstherapeutischen Intervention: untersucht an depressiven Personen. Bachelorthesis an der Goethe-Universität Frankfurt /Main.

Stanton, R. & Reaburn, P. (2014). Exercise and the treatment of depression: A review of the exercise program variables. *Journal of Science and Medicine in Sport*, 17(2), 177–182. <https://doi.org/10.1016/j.jsams.2013.03.010>

Steer, R. A., Rissmiller, D. J. & Beck, A. T. (2000). Use of the Beck Depression Inventory-II with depressed geriatric inpatients. *Behaviour Research and Therapy*, 38(3), 311–318.

Strickland, J. C. & Smith, M. A. (2014). The anxiolytic effects of resistance exercise. *Frontiers in Psychology*, 5, 753. <https://doi.org/10.3389/fpsyg.2014.00753>

Stubbs, B., Firth, J., Berry, A., Schuch, F. B., Rosenbaum, S., Gaughran, F. et al. (2016). How much physical activity do people with schizophrenia engage in? A systematic review, comparative meta-analysis and meta-regression. *Schizophrenia Research*, 176(2–3), 431–440. <https://doi.org/10.1016/j.schres.2016.05.017>

Teychenne, M., Ball, K. & Salmon, J. (2008). Physical activity and likelihood of depression in adults: a review. *Preventive Medicine*, 46(5), 397–411. <https://doi.org/10.1016/j.ypmed.2008.01.009>

Thiel, C., Vogt, L. & Banzer, W. (2011). Bewegung – vielseitige Medizin, die wirkt : dosierte körperliche Aktivität bei chronischen Erkrankungen steigert Gesundheit und Lebensqualität. *Forschung Frankfurt : Wissenschaftsmagazin der Goethe-Universität*, 29(2), 12–19.

Trökes, A. (2017). *Yoga bei Depression*. Freiburg, Breisgau: Herder Verlag.

Vasques, P. E., Moraes, H., Silveira, H., Deslandes, A. C. & Laks, J. (2011). Acute exercise improves cognition in the depressed elderly: the effect of dual-tasks. *Clinics (Sao Paulo, Brazil)*, 66(9), 1553–1557. <https://doi.org/10.1590/s1807-59322011000900008>

Weigelt, M., Steggemann, Y., Machlitt, D. & Engbert, K. (2012). Sport- und Bewegungstherapie bei psychischen Erkrankungen. *PiD - Psychotherapie im Dialog*, 13(04), 91–93. <https://doi.org/10.1055/s-0032-1321414>

WHO Commission on Health and Environment & World Health Organization. (1992). Our planet, our health : report of the WHO Commission on Health and Environment. World Health Organization. <https://apps.who.int/iris/handle/10665/37933>

Wiemeyer, J. & Hänsel, F. (2017). Körperliche Aktivität. In V. Oertel & S. Matura (Hrsg.), *Bewegung und Sport gegen Burnout, Depressionen und Ängste* (S. 3–11). Berlin: Springer. [https://doi.org/10.1007/978-3-662-53938-5\\_1](https://doi.org/10.1007/978-3-662-53938-5_1)

Wipfli, B. M., Rethorst, C. D. & Landers, D. M. (2008). The anxiolytic effects of exercise: a meta-analysis of randomized trials and dose-response analysis. *Journal of Sport & Exercise Psychology*, 30(4), 392–410.

Wittchen, H.-U. & Hoyer, J. (Hrsg.). (2011). *Klinische Psychologie & Psychotherapie* (Springer-Lehrbuch). Berlin: Springer. <https://doi.org/10.1007/978-3-642-13018-2>

Wolf, S. & Hautzinger, M. (2012). Linderung depressiver Symptomatik: Ist sportliche Aktivität eine empfehlenswerte therapeutische Methode? *Psychotherapeut*, 57(3), 241–247. <https://doi.org/10.1007/s00278-011-0854-z>

World Health Organization. What is Moderate-intensity and Vigorous-intensity Physical Activity? [http://www.who.int/dietphysicalactivity/physical\\_activity\\_intensity/en/](http://www.who.int/dietphysicalactivity/physical_activity_intensity/en/)

World Health Organization. (2015). *Physical activity strategy for the WHO European Region 2016–2025*. Copenhagen: World Health Organization.

Wu, T., Gao, X., Chen, M. & van Dam, R. M. (2009). Long-term effectiveness of diet-plus-exercise interventions vs. diet-only interventions for weight loss: a meta-analysis. *Obesity Reviews: An Official Journal of the International Association for the Study of Obesity*, 10(3), 313–323. <https://doi.org/10.1111/j.1467-789X.2008.00547.x>

Yoshitake, T., Kiyohara, Y., Kato, I., Ohmura, T., Iwamoto, H., Nakayama, K. et al. (1995). Incidence and risk factors of vascular dementia and Alzheimer's disease in a defined elderly Japanese population: the Hisayama Study. *Neurology*, 45(6), 1161–1168. <https://doi.org/10.1212/wnl.45.6.1161>